

# 2.5 — OLS: Precision and Diagnostics

ECON 480 • Econometrics • Fall 2021


Ryan Safner

Assistant Professor of Economics

✉ [safner@hood.edu](mailto:safner@hood.edu)

🔗 [ryansafner/metricsF21](https://ryansafner/metricsF21)

🌐 [metricsF21.classes.ryansafner.com](https://metricsF21.classes.ryansafner.com)



# Outline



Variation in  $\hat{\beta}_1$


Presenting Regression Results

Diagnostics about Regression

Problem: Heteroskedasticity

Outliers


# The Sampling Distribution of $\hat{\beta}_1$




$$\hat{\beta}_1 \sim N(E[\hat{\beta}_1], \sigma_{\hat{\beta}_1})$$

1. **Center** of the distribution (last class)

- $E[\hat{\beta}_1] = \beta_1$



# The Sampling Distribution of $\hat{\beta}_1$



$$\hat{\beta}_1 \sim N(E[\hat{\beta}_1], \sigma_{\hat{\beta}_1}^2)$$

1. **Center** of the distribution (last class)


- $E[\hat{\beta}_1] = \beta_1^\dagger$

2. How **precise** is our estimate? (today)

- **Variance**  $\sigma_{\hat{\beta}_1}^2$  or **standard error**<sup>‡</sup>  $\sigma_{\hat{\beta}_1}$

<sup>†</sup> Under the 4 assumptions about  $u$  (particularly,  $cor(X, u) = 0$ ).


<sup>‡</sup> Standard **“error”** is the analog of standard *deviation* when talking about the *sampling distribution* of a sample statistic (such as  $\bar{X}$  or  $\hat{\beta}_1$ ).





# Variation in $\hat{\beta}_1$

# What Affects Variation in $\hat{\beta}_1$




$$\text{var}(\hat{\beta}_1) = \frac{(SER)^2}{n \times \text{var}(X)}$$

$$\text{se}(\hat{\beta}_1) = \sqrt{\text{var}(\hat{\beta}_1)} = \frac{SER}{\sqrt{n} \times \text{sd}(X)}$$

- Variation in  $\hat{\beta}_1$  is affected by 3 things:
  1. **Goodness of fit of the model (SER)<sup>†</sup>**
    - Larger  $SER$  → larger  $\text{var}(\hat{\beta}_1)$
  2. **Sample size,  $n$** 
    - Larger  $n$  → smaller  $\text{var}(\hat{\beta}_1)$
  3. **Variance of  $X$** 
    - Larger  $\text{var}(X)$  → smaller  $\text{var}(\hat{\beta}_1)$


<sup>†</sup> Recall from last class, the **Standard Error of the Regression**  $\hat{\sigma}_u = \sqrt{\frac{\sum \hat{u}_i^2}{n-2}}$

# Variation in $\hat{\beta}_1$ : Goodness of Fit




## Model With Better Fit

Lower SER lowers variation in  $\hat{\beta}_1$




## Model With Worse Fit

Higher SER raises variation in  $\hat{\beta}_1$




# Variation in $\hat{\beta}_1$ : Sample Size




## Model With Fewer Observations

Smaller n raises variation in  $\hat{\beta}_1$




## Model With More Observations

Larger n lowers variation in  $\hat{\beta}_1$






# Variation in $\hat{\beta}_1$ : Variation in $X$




## Model With More Variation in X

Larger  $\text{var}(X)$  lowers variation in  $\hat{\beta}_1$



## Model With Less Variation in X


Smaller  $\text{var}(X)$  raises variation in  $\hat{\beta}_1$





# Presenting Regression Results

# Our Class Size Regression: Base R




- How can we present all of this information in a tidy way?

```
summary(school_reg) # get full summary
```

```
##
## Call:
## lm(formula = testscr ~ str, data = CASchool)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -47.727 -14.251   0.483  12.822  48.540
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  698.9330     9.4675   73.825 < 2e-16 ***
## str          -2.2798     0.4798   -4.751 2.78e-06 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 18.58 on 418 degrees of freedom
## Multiple R-squared:  0.05124,    Adjusted R-squared:  0.04897
## F-statistic: 22.58 on 1 and 418 DF,  p-value: 2.783e-06
```

# Our Class Size Regression: Broom I




- `broom`'s `tidy()` function creates a tidy tibble of regression output

```
# load broom
library(broom)

# tidy regression output
tidy(school_reg)
```

```
## # A tibble: 2 × 5
##   term      estimate std.error statistic  p.value
##   <chr>      <dbl>    <dbl>    <dbl>  <dbl>
## 1 (Intercept) 699.      9.47     73.8 6.57e-242
## 2 str        -2.28     0.480    -4.75 2.78e- 6
```

# Our Class Size Regression: Broom II




- `broom`'s `glance()` gives us summary statistics about the regression

```
glance(school_reg)
```

```
## # A tibble: 1 × 12
##   r.squared adj.r.squared sigma statistic    p.value    df logLik   AIC   BIC
##   <dbl>      <dbl> <dbl>    <dbl>    <dbl> <dbl> <dbl> <dbl> <dbl>
## 1    0.0512      0.0490  18.6     22.6 0.00000278     1 -1822. 3650. 3663.
## # ... with 3 more variables: deviance <dbl>, df.residual <int>, nobs <int>
```

# Presenting Regressions in a Table




- Professional journals and papers often have a **regression table**, including:
  - Estimates of  $\hat{\beta}_0$  and  $\hat{\beta}_1$
  - Standard errors of  $\hat{\beta}_0$  and  $\hat{\beta}_1$  (often below, in parentheses)
  - Indications of statistical significance (often with asterisks)
  - Measures of regression fit:  $R^2$ ,  $SER$ , etc
- Later: multiple rows & columns for multiple variables & models

	Test Score
Intercept	698.93 *** (9.47)
STR	-2.28 *** (0.48)
N	420
R-Squared	0.05
SER	18.58

\*\*\* p < 0.001; \*\* p < 0.01; \* p < 0.05.

# Regression Output with huxtable I




- You will need to first  
`install.packages("huxtable")`
- Load with `library(huxtable)`
- Command: `huxreg()`
- Main argument is the name of your `lm` object
- Default output is fine, but often we want to customize a bit

```
# install.packages("huxtable")  
library(huxtable)  
huxreg(school_reg)
```

	(1)
(Intercept)	698.933 *** (9.467)
str	-2.280 *** (0.480)
N	420
R2	0.051
logLik	-1822.250
AIC	3650.499
*** p < 0.001; ** p < 0.01; * p < 0.05.	

# Regression Output with huxtable II



- Can give title to each column

```
"Test Score" = school_reg
```

- Can change name of coefficients from default

```
coefs = c("Intercept" = "(Intercept)",  
         "STR" = "str")
```

- Decide what statistics to include, and rename them


```
statistics = c("N" = "nobs",  
              "R-Squared" = "r.squared",  
              "SER" = "sigma")
```

- Choose how many decimal places to round to

```
number_format = 2
```



# Regression Output with huxtable III




```
huxreg("Test Score" = school_reg,  
      coefs = c("Intercept" = "(Intercept)",  
               "STR" = "str"),  
      statistics = c("N" = "nobs",  
                   "R-Squared" = "r.squared",  
                   "SER" = "sigma"),  
      number_format = 2)
```

	Test Score
Intercept	698.93 *** (9.47)
STR	-2.28 *** (0.48)
N	420
R-Squared	0.05
SER	18.58

\*\*\* p < 0.001; \*\* p < 0.01; \* p < 0.05.

# Regression Outputs




- `huxtable` is one package you can use
  - See [here for more options](#)
- I used to only use `stargazer`, but as it was originally meant for STATA, it has limits and problems
  - A great [cheatsheet](#) by my friend Jake Russ




# Diagnostics about Regression

# Diagnostics: Residuals I



- We often look at the residuals of a regression to get more insight about its **goodness of fit** and its **bias**
- Recall `broom`'s `augment` creates some useful new variables
  - `.fitted` are fitted (predicted) values from model, i.e.  $\hat{Y}_i$
  - `.resid` are residuals (errors) from model, i.e.  $\hat{u}_i$

# Diagnostics: Residuals II




- Often a good idea to store in a new object (so we can make some plots)

```
aug_reg<-augment(school_reg)
```

```
aug_reg %>% head()
```

<b>testscr</b>	<b>str</b>	<b>.fitted</b>	<b>.resid</b>	<b>.hat</b>	<b>.sigma</b>	<b>.cooksd</b>	<b>.std.resid</b>
691	17.9	658	32.7	0.00442	18.5	0.00689	1.76
661	21.5	650	11.3	0.00475	18.6	0.000893	0.612
644	18.7	656	-12.7	0.00297	18.6	0.0007	-0.685
648	17.4	659	-11.7	0.00586	18.6	0.00117	-0.629
641	18.7	656	-15.5	0.00301	18.6	0.00105	-0.836
606	21.4	650	-44.6	0.00446	18.5	0.013	-2.4

# Recap: Assumptions about Errors



- We make **4 critical assumptions about  $u$** :

1. The expected value of the residuals is 0

$$E[u] = 0$$

2. The variance of the residuals over  $X$  is constant:

$$\text{var}(u|X) = \sigma_u^2$$

3. Errors are not correlated across observations:


$$\text{cor}(u_i, u_j) = 0 \quad \forall i \neq j$$

4. There is no correlation between  $X$  and the error term:

$$\text{cor}(X, u) = 0 \text{ or } E[u|X] = 0$$



# Assumptions 1 and 2: Errors are i.i.d.




- Assumptions 1 and 2 assume that errors are coming from the same (*normal*) distribution


$$u \sim N(0, \sigma_u)$$

- Assumption 1:  $E[u] = 0$
- Assumption 2:  $sd(u|X) = \sigma_u$ 
  - virtually always unknown...
- We often can visually check by plotting a **histogram** of  $u$

# Plotting Residuals




```
ggplot(data = aug_reg)+  
  aes(x = .resid)+  
  geom_histogram(color="white", fill = "pink")+  
  labs(x = expression(paste("Residual, ", hat(u))))+  
  theme_pander(base_family = "Fira Sans Condensed",  
              base_size=20)
```





# Plotting Residuals




```
ggplot(data = aug_reg)+  
  aes(x = .resid)+  
  geom_histogram(color="white", fill = "pink")+  
  labs(x = expression(paste("Residual, ", hat(u))))+  
  theme_pander(base_family = "Fira Sans Condensed",  
              base_size=20)
```


- Just to check:

```
aug_reg %>%  
  summarize(E_u = mean(.resid),  
            sd_u = sd(.resid))
```

<b>E_u</b>	<b>sd_u</b>
3.7e-13	18.6




# Residual Plot



- We often plot a **residual plot** to see any odd patterns about residuals
  - $x$ -axis are  $X$  values (`str`)
  - $y$ -axis are  $u$  values (`.resid`)


```
ggplot(data = aug_reg)+  
  aes(x = str,  
      y = .resid)+  
  geom_point(color="blue")+  
  geom_hline(aes(yintercept = 0), color="red")+  
  labs(x = "Student to Teacher Ratio",  
       y = expression(paste("Residual, ", hat(u))))  
  theme_pander(base_family = "Fira Sans Condensed",  
              base_size=20)
```





# Problem: Heteroskedasticity

# Homoskedasticity




- "**Homoskedasticity**:" variance of the residuals over  $X$  is constant, written:

$$\text{var}(u|X) = \sigma_u^2$$

- Knowing the value of  $X$  does not affect the variance (spread) of the errors



# Heteroskedasticity I




- "**Heteroskedasticity**:" variance of the residuals over  $X$  is *NOT* constant:

$$\text{var}(u|X) \neq \sigma_u^2$$

- **This does not cause  $\hat{\beta}_1$  to be biased**, but it does cause the standard error of  $\hat{\beta}_1$  to be incorrect
- This **does** cause a problem for **inference**!



# Heteroskedasticity II




- Recall the formula for the standard error of  $\hat{\beta}_1$ :

$$se(\hat{\beta}_1) = \sqrt{\text{var}(\hat{\beta}_1)} = \frac{SER}{\sqrt{n} \times sd(X)}$$

- This actually *assumes* homoskedasticity

# Heteroskedasticity III




- Under heteroskedasticity, the standard error of  $\hat{\beta}_1$  mutates to:


$$se(\hat{\beta}_1) = \sqrt{\frac{\sum_{i=1}^n (X_i - \bar{X})^2 \hat{u}^2}{\left[ \sum_{i=1}^n (X_i - \bar{X})^2 \right]^2}}$$

- This is a **heteroskedasticity-robust** (or just "**robust**") method of calculating  $se(\hat{\beta}_1)$
- Don't learn formula, **do learn what heteroskedasticity is and how it affects our model!**

# Visualizing Heteroskedasticity I




- Our original scatterplot with regression line







# Visualizing Heteroskedasticity I




- Our original scatterplot with regression line
- Does the spread of the errors change over different values of  $str$ ?
  - No: homoskedastic
  - Yes: heteroskedastic




# Visualizing Heteroskedasticity I




- Our original scatterplot with regression line
- Does the spread of the errors change over different values of  $str$ ?
  - No: homoskedastic
  - Yes: heteroskedastic




# Heteroskedasticity: Another View




- Using the `ggridges` package
- Plotting the (conditional) distribution of errors by STR
- See that the variation in errors ( $\hat{u}$ ) changes across class sizes!




# More Obvious Heteroskedasticity





- Visual cue: data is "fan-shaped"
  - Data points are closer to line in some areas
  - Data points are more spread from line in other areas




# More Obvious Heteroskedasticity




- Visual cue: data is "fan-shaped"
  - Data points are closer to line in some areas
  - Data points are more spread from line in other areas




# Heteroskedasticity: Another View



- Using the `ggribes` package
- Plotting the (conditional) distribution of errors by  $x$




# What Might Cause Heteroskedastic Errors?




$$\widehat{wage}_i = \hat{\beta}_0 + \hat{\beta}_1 educ_i$$

	Wage
Intercept	-0.90 (0.68)
Years of Schooling	0.54 *** (0.05)
N	526
R-Squared	0.16
SER	3.38

\*\*\* p < 0.001; \*\* p < 0.01; \* p < 0.05.





# What Might Cause Heteroskedastic Errors?



$$\widehat{wage}_i = \hat{\beta}_0 + \hat{\beta}_1 educ_i$$


	Wage
Intercept	-0.90 (0.68)
Years of Schooling	0.54 *** (0.05)
N	526
R-Squared	0.16
SER	3.38

\*\*\*  $p < 0.001$ ; \*\*  $p < 0.01$ ; \*  $p < 0.05$ .







# Heteroskedasticity: Another View



- Using the `ggridges` package
- Plotting the (conditional) distribution of errors by education



# Detecting Heteroskedasticity I




- Several tests to check if data is heteroskedastic
- One common test is **Breusch-Pagan test**
- Can use `bptest()` with `lmtest` package in `R`
  - $H_0$ : homoskedastic
  - If  $p\text{-value} < 0.05$ , reject  $H_0 \implies$  heteroskedastic

```
# install.packages("lmtest")  
library("lmtest")  
bptest(school_reg)
```

```
##  
##      studentized Breusch-Pagan test  
##  
## data:  school_reg  
## BP = 5.7936, df = 1, p-value = 0.01608
```

# Detecting Heteroskedasticity II




- How about our wage regression?

```
# install.packages("lmtest")  
library("lmtest")  
bptest(wage_reg)
```

```
##  
##      studentized Breusch-Pagan test  
##  
## data:  wage_reg  
## BP = 15.306, df = 1, p-value = 9.144e-05
```

# Fixing Heteroskedasticity I




- Heteroskedasticity is easy to fix with software that can calculate **robust** standard errors (using the more complicated formula above)
- Easiest method is to use `estimatr` package
  - `lm_robust()` command (instead of `lm`) to run regression
  - set `se_type="stata"` to calculate robust SEs using the formula above

```
#install.packages("estimatr")  
library(estimatr)  
  
school_reg_robust <-lm_robust(testscr ~ str, data = CASchool,  
                             se_type = "stata")  
  
school_reg_robust
```

```
##           Estimate Std. Error   t value      Pr(>|t|)   CI Lower   CI Upper  
## (Intercept) 698.932952 10.3643599 67.436191 9.486678e-227 678.560192 719.305713  
## str         -2.279808  0.5194892 -4.388557 1.446737e-05 -3.300945 -1.258671  
##           DF
```

# Fixing Heteroskedasticity II




```
library(huxtable)
huxreg("Normal" = school_reg,
      "Robust" = school_reg_robust,
      coefs = c("Intercept" = "(Intercept)",
               "STR" = "str"),
      statistics = c("N" = "nobs",
                    "R-Squared" = "r.squared",
                    "SER" = "sigma"),
      number_format = 2)
```

	Normal	Robust
Intercept	698.93 ***	698.93 ***
	(9.47)	(10.36)
STR	-2.28 ***	-2.28 ***
	(0.48)	(0.52)
N	420	420
R-Squared	0.05	0.05
SER	18.58	

\*\*\* p < 0.001; \*\* p < 0.01; \* p < 0.05.

# Assumption 3: No Serial Correlation



- Errors are not correlated across observations:

$$\text{cor}(u_i, u_j) = 0 \quad \forall i \neq j$$


- For simple cross-sectional data, this is rarely an issue
- Time-series & panel data nearly always contain **serial correlation** or **autocorrelation** between errors
- Errors may be **clustered**
  - **by group**: e.g. all observations from Maryland, all observations from Virginia, etc.
  - **by time**: GDP in 2006 around the world, GDP in 2008 around the world, etc.






# Outliers

# Outliers Can Bias OLS! I




- **Outliers** can affect the slope (and intercept) of the line and add **bias**
  - May be result of human error (measurement, transcribing, etc)
  - May be meaningful and accurate
- In any case, compare how including/dropping outliers affects regression and always discuss outliers!






# Outliers Can Bias OLS! II




```
huxreg("No Outliers" = school_reg,  
      "Outliers" = school_outlier_reg,  
      coefs = c("Intercept" = "(Intercept)",  
               "STR" = "str"),  
      statistics = c("N" = "nobs",  
                   "R-Squared" = "r.squared",  
                   "SER" = "sigma"),  
      number_format = 2)
```

	No Outliers	Outliers
Intercept	698.93 *** (9.47)	641.40 *** (11.21)
STR	-2.28 *** (0.48)	0.71 (0.57)
N	420	423
R-Squared	0.05	0.00
SER	18.58	23.76



# Detecting Outliers



- The `car` package has an `outlierTest` command to run on the regression

```
library("car")
```

```
# Use Bonferonni test
```

```
outlierTest(school_outlier_reg) # will point out which obs #s seem outliers
```

```
##      rstudent unadjusted p-value Bonferroni p
## 422 8.822768      3.0261e-17  1.2800e-14
## 423 7.233470      2.2493e-12  9.5147e-10
## 421 6.232045      1.1209e-09  4.7414e-07
```

```
# find these observations
```

```
CA.outlier %>%
  slice(c(422,423,421))
```

observat	district	testscr	str
422	Crazy School 2	850	28
423	Crazy School 3	820	29
421	Crazy School 1	800	30